

A mu-differentiable Lagrange multiplier rule*

Ricardo Almeida and Delfim F. M. Torres

Department of Mathematics

University of Aveiro

3810-193 Aveiro, Portugal

{ricardo.almeida, delfim}@ua.pt

ABSTRACT

We present some properties of the gradient of a mu-differentiable function. The Method of Lagrange Multipliers for mu-differentiable functions is then exemplified.

Keywords: nonstandard analysis, mu-differentiability, Method of Lagrange Multipliers.

2000 Mathematics Subject Classification: 26E35, 26E05, 26B05.

1 Introduction

In [1] we introduce a new kind of differentiation, what we call mu-differentiability, and we prove necessary and sufficient conditions for the existence of extrema points. For the necessary background on Nonstandard Analysis and for notation, we refer the reader to [1] and references therein. Here we just recall the necessary results.

Definition 1.1. [1] Given an internal function $f : {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}$, we say that $\alpha \in \mathbb{R}^n$ is a *local m-minimum* of f if

$$f(x) \gtrsim f(\alpha) \text{ for all } x \in {}^*B_r(\alpha),$$

where $r \in \mathbb{R}$ is a positive real number.

The crucial fact is that there exists a relationship between m-minimums and minimums:

Lemma 1.1. [1] If $f : {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}$ is mu-differentiable, then

α is a m-minimum of f if and only if α is a minimum of $st(f)$.

With this lemma, and using the fact that

$$st\left(\frac{\partial f}{\partial x_i}\Big|_{\alpha}\right) = \frac{\partial st(f)}{\partial x_i}\Big|_{\alpha} \text{ for } i \in \{1, \dots, n\}, \quad (1.1)$$

it follows:

*Supported by *Centre for Research on Optimization and Control* (CEOC) from the “Fundação para a Ciência e a Tecnologia” (FCT), cofinanced by the European Community Fund FEDER/POCI 2010. Accepted (22/June/2008) for International Journal of Mathematics and Statistics (IJMS), Vol. 4, No. S09, Spring 2009 (in press).

Theorem 1.2. [1] If $f : {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}$ is a mu-differentiable function and α is a m -minimum of f , then

$$\left. \frac{\partial f}{\partial x_i} \right|_{\alpha} \approx 0, \text{ for every } i = 1, \dots, n.$$

In this paper we develop further the theory initiated in [1], proving some properties of the gradient vector (section 2) and a Method of Lagrange Multipliers (section 3). Illustrative examples show the analogy with the classical case.

2 The Gradient Vector

In the sequel f denotes an internal mu-differentiable function from ${}^*\mathbb{R}^n$ to ${}^*\mathbb{R}$.

Definition 2.1. A *gradient vector* of f at $x \in ns({}^*\mathbb{R}^n)$ is defined by

$$\nabla f(x) := \left(\left. \frac{\partial f}{\partial x_1} \right|_x, \dots, \left. \frac{\partial f}{\partial x_n} \right|_x \right)$$

where

$$\left. \frac{\partial f}{\partial x_i} \right|_x \approx \frac{f(x_1, \dots, x_{i-1}, x_i + \epsilon, x_{i+1}, \dots, x_n) - f(x_1, \dots, x_n)}{\epsilon}$$

and ϵ is an infinitesimal satisfying $|\epsilon| > \delta_f$.

Remark 2.1. The positive infinitesimal δ_f that appears in Definition 2.1 is given by the m -differentiability of f (cf. [1]).

Remark 2.2. Observe that

$$\left. \frac{\partial f}{\partial x_i} \right|_x \approx Df_x(e_i),$$

where $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ denotes the i th canonical vector, and Df_x denotes the derivative operator of f at x .

Theorem 2.1. If $x, y \in ns({}^*\mathbb{R}^n)$ and $x \approx y$, then

$$\left. \frac{\partial f}{\partial x_i} \right|_x \approx \left. \frac{\partial f}{\partial x_i} \right|_y, \quad i = 1, \dots, n,$$

i.e., $\nabla f(x) \approx \nabla f(y)$.

Proof. Simply observe that $Df_x(e_i) \approx Df_y(e_i)$. □

Theorem 2.2. If $u \in {}^*\mathbb{R}^n$ is a finite vector, then

$$\forall x \in ns({}^*\mathbb{R}^n) \quad Df_x(u) \approx \nabla f(x) \cdot u.$$

Proof. Since $st(f)$ is a C^1 function, it follows that for any $v \in \mathbb{R}^n$

$$Dst(f)_{st(x)}(v) = \nabla st(f)(st(x)) \cdot v.$$

By the Transfer Principle of Nonstandard Analysis, it still holds for $u \in {}^*\mathbb{R}^n$. On the other hand,

$$1. \quad Dst(f)_{st(x)}(v) = st(Df_{st(x)})(u) \approx Df_x(u),$$

$$2. \nabla st(f)(st(x)) = st(\nabla f(st(x))) \approx \nabla f(x),$$

which proves the desired. $\square$

We point out that, in opposite to classical functions, if $\nabla f(x)$ is a gradient vector of f at x , then $\nabla f(x) + \Omega$, where $\Omega \in {}^*\mathbb{R}^n$ is an infinitesimal vector, is also a gradient vector at x . Conversely, if $\nabla f(x)$ and $\nabla^1 f(x)$ are two gradient vectors, then $\nabla f(x) - \nabla^1 f(x) \approx 0$.

From now on, when there is no danger of confusion, we simply write ∇f instead of $\nabla f(x)$.

Example 2.1. Let $f(x, y, z) = (1+\epsilon)xy^2 - \delta z$, with $(x, y, z) \in {}^*\mathbb{R}^3$, and ϵ and δ be two infinitesimal numbers. Given an infinitesimal θ ,

$$\begin{aligned} \frac{(1+\epsilon)(x+\theta)y^2 - \delta z - ((1+\epsilon)xy^2 - \delta z)}{\theta} &= (1+\epsilon)y^2, \\ \frac{(1+\epsilon)x(y+\theta)^2 - \delta z - ((1+\epsilon)xy^2 - \delta z)}{\theta} &= 2(1+\epsilon)xy + \theta(1+\epsilon)x, \\ \frac{(1+\epsilon)xy^2 - \delta(z+\theta) - ((1+\epsilon)xy^2 - \delta z)}{\theta} &= -\delta, \end{aligned}$$

and we can choose

$$\frac{\partial f}{\partial x} = (1+\epsilon)y^2, \quad \frac{\partial f}{\partial y} = 2(1+\epsilon)xy, \quad \frac{\partial f}{\partial z} = -\delta.$$

Theorem 2.3. *If f and g are mu-differentiable and $k \in fin({}^*\mathbb{R})$, then*

$$\nabla(kf) = k\nabla f, \quad \nabla(f+g) = \nabla f + \nabla g, \quad \text{and} \quad \nabla(fg) = f\nabla g + g\nabla f.$$

Proof. We prove only the last equality. Fix an infinitesimal number ϵ such that $|\epsilon| > \delta_f$. Then,

$$\begin{aligned} \frac{\partial(fg)}{\partial x_i} &\approx \frac{(fg)(x_1, \dots, x_{i-1}, x_i + \epsilon, x_{i+1}, \dots, x_n) - (fg)(x_1, \dots, x_n)}{\epsilon} \\ &= f(x_1, \dots, x_n) \frac{g(x_1, \dots, x_{i-1}, x_i + \epsilon, x_{i+1}, \dots, x_n) - g(x_1, \dots, x_n)}{\epsilon} \\ &\quad + g(x_1, \dots, x_{i-1}, x_i + \epsilon, x_{i+1}, \dots, x_n) \frac{f(x_1, \dots, x_{i-1}, x_i + \epsilon, x_{i+1}, \dots, x_n) - f(x_1, \dots, x_n)}{\epsilon} \\ &\approx f(x) \frac{\partial g}{\partial x_i} + g(x) \frac{\partial f}{\partial x_i} \end{aligned}$$

by the continuity of g . $\square$

Definition 2.2. We say that x is a *m-critical point* of f if $\nabla f(x) \approx 0$.

The following lemma is an immediate consequence of (1.1) and Definition 2.2.

Lemma 2.4. *A point x is a m-critical point of f if and only if $st(x)$ is a critical point of $st(f)$.*

3 The Method of Lagrange Multipliers

Let $f : {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}$ and $g_j : {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}$, $j = 1, \dots, m$ ($m \in \mathbb{N}$, $m < n$), denote internal mu-differentiable functions. We address the problem of finding m-minimums or m-maximums of f , subject to the conditions $g_j(x) \approx 0$, for all j . The constraints $g_j(x) \approx 0$, $j = 1, \dots, m$, are called *side conditions*. Lagrange solved this problem (for standard differentiable functions), introducing new variables, $\lambda_1, \dots, \lambda_m$, and forming the augmented function

$$F(x, \lambda_1, \dots, \lambda_m) = f(x) + \sum_{j=1}^m \lambda_j g_j(x), \quad x \in \mathbb{R}^n.$$

Roughly speaking, Lagrange proved that the problem of finding the critical points of f , satisfying the conditions $g_j(x) = 0$, is equivalent to find the critical points of F . We present here a method to determine critical points for internal functions with side conditions, based on the *Method of Lagrange Multipliers*. Similarly to the classical setting, define

$$\begin{aligned} F : \quad {}^*\mathbb{R}^{n+m} &\rightarrow {}^*\mathbb{R} \\ (x_1, \dots, x_n, \lambda_1, \dots, \lambda_m) &\mapsto f(x_1, \dots, x_n) + \sum_{j=1}^m \lambda_j g_j(x_1, \dots, x_n). \end{aligned}$$

If we let $g := (g_1, \dots, g_m)$ and $\lambda := (\lambda_1, \dots, \lambda_m)$, we can simply write

$$F(x, \lambda) = f(x) + \lambda \cdot g(x). \quad (3.1)$$

Theorem 3.1. [Lagrange rule in normal form with one constraint] Let $f : {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}$ and $g : {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}$ be two mu-differentiable functions, and α a m-minimum of f such that $g(\alpha) \approx 0$ and $\nabla g(\alpha) \not\approx 0$. Then, there exists a finite $\lambda \in {}^*\mathbb{R}$ such that

$$\nabla f(\alpha) + \lambda \nabla g(\alpha) \approx 0.$$

Proof. Since $st(f)$ and $st(g)$ are functions of class C^1 , α is a minimum of $st(f)$, $st(g)(\alpha) = 0$ and $\nabla st(g)(\alpha) \neq 0$. It follows (see, e.g., [2, p. 148]) that

$$\exists \lambda \in \mathbb{R} \quad \nabla st(f)(\alpha) + \lambda \nabla st(g)(\alpha) = 0.$$

Hence,

$$\nabla f(\alpha) + \lambda \nabla g(\alpha) \approx 0.$$

□

Remark 3.1. Suppose that we are in the conditions of Theorem 3.1. Then, there exists some $\lambda_1 \in fin({}^*\mathbb{R})$ such that

$$\nabla f(\alpha) + \lambda_1 \nabla g(\alpha) \approx 0,$$

i.e.,

$$\left. \frac{\partial f}{\partial x_i} \right|_{\alpha} + \lambda_1 \left. \frac{\partial g}{\partial x_i} \right|_{\alpha} \approx 0, \quad i = 1, \dots, n.$$

Using the notation (3.1), if α is a m-minimum of f and $g(\alpha) \approx 0$, then

$$\begin{cases} \left. \frac{\partial F}{\partial x_i} \right|_{(\alpha, \lambda_1)} = \left. \frac{\partial f}{\partial x_i} \right|_{\alpha} + \lambda_1 \left. \frac{\partial g}{\partial x_i} \right|_{\alpha} \approx 0, & i = 1, \dots, n, \\ \left. \frac{\partial F}{\partial \lambda} \right|_{(\alpha, \lambda_1)} \approx g(\alpha) \approx 0. \end{cases} \quad (3.2)$$

Consequently, the m-critical points are solutions of the system

$$\frac{\partial F}{\partial x_i} \approx 0, i = 1, \dots, n, \text{ and } \frac{\partial F}{\partial \lambda} \approx 0,$$

i.e., $\nabla F \approx 0$.

Example 3.1. Let $f(x, y, z) = xyz + \epsilon$, with $\epsilon \approx 0$, and consider the constraint $g(x, y, z) = x^2 + 2(y + \delta)^2 + 3z^2 - 1$, with $\delta \approx 0$. In this case, we define

$$F(x, y, z, \lambda) := xyz + \epsilon + \lambda(x^2 + 2(y + \delta)^2 + 3z^2 - 1).$$

The system (3.2) takes the form

$$\begin{cases} yz + 2\lambda x \approx 0 \\ xz + 4\lambda(y + \delta) \approx 0 \\ xy + 6\lambda z \approx 0 \\ x^2 + 2(y + \delta)^2 + 3z^2 - 1 \approx 0. \end{cases}$$

Since

$$xyz \approx -2\lambda x^2 \approx -4\lambda y(y + \delta) \approx -6\lambda z^2,$$

if $\lambda \not\approx 0$, the solution is

$$x^2 \approx \frac{1}{3}, y^2 \approx \frac{1}{6} \text{ and } z^2 \approx \frac{1}{9};$$

if $\lambda \approx 0$, then

$$\left(0, 0, \pm \frac{1}{\sqrt{3}}\right), \left(0, \pm \frac{1}{\sqrt{2}}, 0\right) \text{ and } (\pm 1, 0, 0)$$

are solutions. Observe that

$$\nabla g = (2x, 4(y + \delta), 6z) \approx (0, 0, 0) \text{ if and only if } (x, y, z) \approx (0, 0, 0).$$

One easily checks that

$$f\left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{6}}, \frac{1}{3}\right) = \frac{1}{3\sqrt{18}} + \epsilon \text{ is the m-maximum and}$$

$$f\left(-\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{6}}, \frac{1}{3}\right) = -\frac{1}{3\sqrt{18}} + \epsilon \text{ is the m-minimum}$$

of f subject to the constraint g .

We now prove a more general Lagrange rule, admitting possibility of abnormal critical points ($\mu = 0$) and multiple constraints.

Theorem 3.2 (Lagrange rule). *Let $f, g_1, \dots, g_m$ be mu-differentiable functions on ${}^*\mathbb{R}^n$. Let α be a m-minimum of f satisfying*

$$g_1(\alpha) \approx \dots \approx g_m(\alpha) \approx 0.$$

Then, there exist finite hyper-reals $\mu, \lambda_1, \dots, \lambda_m \in {}^\mathbb{R}$, not all infinitesimals, such that*

$$\mu \nabla f(\alpha) + \lambda_1 \nabla g_1(\alpha) + \dots + \lambda_m \nabla g_m(\alpha) \approx 0.$$

Remark 3.2. Defining $F(x, \mu, \lambda) := \mu f(x) + \lambda \cdot g(x)$, the necessary optimality condition given by Theorem 3.2 can be written as $\partial F/\partial x \approx \partial F/\partial \lambda \approx 0$.

Proof. First observe that $st(f), st(g_1), \dots, st(g_m)$ are all functions of class C^1 , $\nabla st(f)(\alpha) = st(\nabla f)(\alpha)$ and $\nabla st(g_j)(\alpha) = st(\nabla g_j)(\alpha)$, for $j = 1, \dots, m$. Furthermore, since α is a minimum of $st(f)$ and

$$st(g_1)(\alpha) = \dots = st(g_m)(\alpha) = 0,$$

there exist reals $\mu, \lambda_1, \dots, \lambda_m$, not all zero, such that

$$\mu \nabla st(f)(\alpha) + \lambda_1 \nabla st(g_1)(\alpha) + \dots + \lambda_m \nabla st(g_m)(\alpha) = 0$$

(see, e.g., [2, p. 148]). Consequently,

$$\mu st(\nabla f)(\alpha) + \lambda_1 st(\nabla g_1)(\alpha) + \dots + \lambda_m st(\nabla g_m)(\alpha) = 0. \quad (3.3)$$

On the other hand, we have

$$\mu st(\nabla f)(\alpha) = \mu st(\nabla f(\alpha)) \approx \mu \nabla f(\alpha).$$

Analogously, for each $j = 1, \dots, m$,

$$\lambda_j st(\nabla g_j)(\alpha) \approx \lambda_j \nabla g_j(\alpha).$$

Substituting on equation (3.3) the previous relations, one proves the desired result. $\square$

Example 3.2. Let $f(x, y, z) = z^2/2 - (x + \epsilon)y$, with $\epsilon \approx 0$, be the function to be extremized, and $g_1(x, y, z) = x^2 + y - 1$ and $g_2(x, y, z) = x + z - 1 + \delta$, with $\delta \approx 0$, be the constraints. Then, the augmented function is

$$F(x, y, z, \mu, \lambda_1, \lambda_2) = \mu [z^2/2 - (x + \epsilon)y] + \lambda_1(x^2 + y - 1) + \lambda_2(x + z - 1 + \delta).$$

To find the local extrema of f , subject to the conditions $g_1 \approx 0$ and $g_2 \approx 0$, we form the system

$$\begin{cases} -\mu y + 2\lambda_1 x + \lambda_2 \approx 0 \\ -\mu(x + \epsilon) + \lambda_1 \approx 0 \\ \mu z + \lambda_2 \approx 0 \\ x^2 + y - 1 \approx 0 \\ x + z - 1 + \delta \approx 0 \end{cases} \quad (3.4)$$

of necessary optimality conditions. Assume $\mu \approx 0$ (abnormal case). Then, the first two equations in (3.4) imply immediately that $\lambda_1 \approx \lambda_2 \approx 0$. This is not a possibility by Theorem 3.2. We conclude that $\mu \not\approx 0$. The solutions of (3.4) are then infinitely close to the vectors

$$(-1, 0, 2) \quad \text{and} \quad (2/3, 5/9, 1/3).$$

Hence, if f has any m-extrema under the given constraints, then they must occur at either $(-1, 0, 2)$ or $(2/3, 5/9, 1/3)$.

References

- [1] R. Almeida and D. F. M. Torres. Relaxed optimality conditions for mu-differentiable functions. *Int. J. Appl. Math. Stat.* (accepted). [arXiv:0806.3545v1 \[math.CA\]](#)
- [2] W. Cheney. *Analysis for Applied Mathematics*. GTM. Springer-Verlag, New York, 2001.